

Exercise - Week 6

- ① Show that the forgetful functor $U: \text{Mon} \rightarrow \text{Set}$ does not have a right adjoint. What about groups/rings?
- ② The free functor $F: \text{Set} \rightarrow \text{Mon}$ sending a set to the list monoid is left adjoint to $U: \text{Mon} \rightarrow \text{Set}$. But does F have a left adjoint?
- ③ Consider the forgetful functor $U: \text{Grph} \rightarrow \text{Set}$ from graphs to sets.
- need Graph to mean graph with loops, if we want $\mathbb{I}\mathbb{I}\mathbb{I}$!
- Show that there are adjoint functors $\mathbb{I}\mathbb{I}\dashv F \dashv U \dashv R$ and that this string of adjunctions cannot be extended any further.
- ④ Consider adjoint functors
- $$\mathcal{A} \xrightleftharpoons[\mathbf{U}_1]{F_1} \mathcal{B} \xrightleftharpoons[\mathbf{U}_2]{F_2} \mathcal{C}.$$
- Show that we have an adjunction $F_1 F_2 \dashv U_2 U_1$.

Limits as adjoints

- let J be a small cat & \mathcal{C} a category.

- Given $a \in \mathcal{C}$ we can define the constant functor $\Delta_a: J \rightarrow \mathcal{C} : j \mapsto a$
at a $j \xrightarrow{a} k \mapsto a \xrightarrow{\text{id}} a$

- Show that a natural transformation $\Delta_a \xrightarrow{f} D$ is the same thing as a cone $(A \xrightarrow{f_i} D_i)_{i \in J}$.

- Show that Δ defines a functor

$$\mathcal{C} \xrightarrow{\Delta} [J, \mathcal{C}] \sim \text{Functor cat}$$

& that Δ has a left adj. \Leftrightarrow
 \mathcal{C} has all colimits of shape J .