



Lead Author e-mail: mbartak@sci.muni.cz

Title: *Photosynthetic characteristics of lichens of genus Umbilicaria from SW Greenland (Nuuk area) in response to thallus dehydration*

Bartak, M.¹, Hansen, E.S.², Hazdrova, J.¹, Skacelova, K.¹, Trnkova, K.¹, Prosek, P.³

¹*Department of Experimental Biology, Masaryk University, Kamenice 5, Brno, Czech Republic*

²*Natural History Museum of Denmark, Gothersgade 130, 1123 København, Denmark*

³*Department of Geography, Faculty of Science, Masaryk University, Kotlářská 2, Brno, Czech Republic*

Thalli of foliose epilithic lichens *Umbilicaria arctica* and *U.hyperborea* were collected on the rocks at several locations in the neighbourhood of Nuuk, and transported to the Czech Republic where kept in dry state in dark at 5 °C before experiments. After 48 h rehydration, simultaneous measurements of (1)effective quantum yield (Yield_{PSII}) of photosystem II, (2)photochemical reflectance index (PRI), (3)normalized difference vegetation index (NDVI), (4)chlorophyll fluorescence fast kinetics (OJIP) were made in response to gradual thallus dehydration expressed as water potential (WP). Dehydration-response curves of Yield_{PSII} showed S-curve relationship. In both species, the first signs of inhibition of photosynthetic processes appeared at WP of about -10 MPa. Further dehydration led to a decrease in Yield_{PSII} and, finally, full inhibition of PSII photochemical photosynthetic processes. Critical point for PSII processes was found at WP of about -25 MPa, similarly to other epilithic lichens investigated by this method (Bartak et al. 2005). In our study, *U.arctica* showed higher Yield_{PSII} and less sensitivity to dehydration than *U.hyperborea* in the WP range of -5 to -15 MPa. Similarly to previous study (Jupa et al. 2012) done on *U.cylindrica* and *U.decussata* from Svalbard, PRI in *U.arctica* and *U.hyperborea* exhibited curvilinear increase with dehydration. The relation of PRI to WP was, however, species-specific. NDVI, a vigor indicator, decreased with dehydration in both species, however, due to generally black color of *U. hyperborea*, the decrease was much less pronounced in the species. OJIPS recorded in fully hydrated thalli showed typical polyphasic curves with peak chlorophyll fluorescence level "P" found at 150-300 ms followed by a dip "D" typical for trebouxoid lichens (Ilik et al 2006). With more pronounced dehydration, OJIPs exhibited a decrease in chlorophyll fluorescence signal and photosynthetic parameters derived from OJIPs. These changes indicated dehydration-dependent inhibition of photosynthetic processes in both studied species. All the results presented in this study indicated a high degree of tolerance of the two *Umbilicaria* species to partial dehydration stress. However, interspecific differences in photosynthetic parameters were apparent in response to thalli dehydration. **Acknowledgements:** The authors are grateful to the CzechPolar infrastructure for providing laboratory facilities necessary for the above experiments.

References:



Bartak, M. et al. (2005): *The Lichenologist*, 37: 433-443

Ilik, P. et al. (2006): *Biochimica et Biophysica Acta - Bioenergetics*. 1757: 12-20.

Jupa, R. et al. (2012): *Czech Polar Reports*, 2: 31-41.