



# The mediating role of air pollutants in the association between education and lung function among the elderly, the HAPIEE study

Consuelo Quispe-Haro<sup>a</sup>, Daniel Szabó<sup>a</sup>, Katarzyna Kordas<sup>b</sup>, Nadezda Capkova<sup>c</sup>, Hynek Pikhart<sup>a,d,\*</sup>, Martin Bobak<sup>a,d</sup>

<sup>a</sup> RECETOX, Faculty of Science, Masaryk University, Kotlarska 2, Brno, Czech Republic

<sup>b</sup> Department of Epidemiology and Environmental Health, University at Buffalo, Buffalo, NY, USA

<sup>c</sup> National Institute of Public Health, Prague, Czech Republic

<sup>d</sup> Research Department of Epidemiology and Public Health, University College London, London, UK


## HIGHLIGHTS

- We investigated lung function in older adults from seven Czech cities.
- We noticed an educational gradient in lung function and exposure to air pollutants.
- PM<sub>10</sub> and NO<sub>2</sub> partly mediated the association between education and lung function.
- Educational attainment had direct and indirect association with lung function.

## GRAPHICAL ABSTRACT

### Mediation effects of PM<sub>10</sub> and NO<sub>2</sub> between education and lung function.

Path analysis revealed a direct positive effect of education on FEV1, while about 12% of the relationship between education and lung function was mediated by PM<sub>10</sub> and NO<sub>2</sub>.



Higher levels of participants' education were associated with lower exposures to PM<sub>10</sub> and NO<sub>2</sub> in their place of residence. Higher exposure to PM<sub>10</sub> was associated with lower lung function.

## ARTICLE INFO

Editor: Lidia Minguéz Alarcon

### Keywords:

Socioeconomic factors  
Air pollution  
Lung function  
Respiratory health  
Environmental health

## ABSTRACT

**Background:** Chronic exposure to air pollutants harms human health, and at a geographical level, concentrations of air pollutants are often associated with socioeconomic disadvantage.

**Objectives:** The aim of this study was to investigate the effects of educational attainment and air pollution on lung function in older adults, and whether air pollution may mediate the effect of education.

**Methods:** The study included 6381 individuals (mean age 58.24 ± 7.14 years) who participated in the Czech HAPIEE (Health, Alcohol, and Psychosocial Factors in Eastern Europe) study. Participants' residential addresses were linked to air pollution data, including mean exposures to PM<sub>10</sub> (particulate matter of aerodynamic diameter below 10 μm) and NO<sub>2</sub> (nitrogen dioxide). We used path analysis to link educational attainment and air pollutants to a standardized measure of the Forced Expiratory Volume in the first second (FEV1).

**Results:** Higher levels of participants' education were associated with lower exposures to PM<sub>10</sub> and NO<sub>2</sub>. Individuals with tertiary education had higher standardized FEV1 than individuals with primary education (88 % vs 95 %). Path analysis revealed a direct positive effect of education on FEV1, while about 12 % of the relationship between education and lung function was mediated by PM<sub>10</sub> and NO<sub>2</sub>.

**Conclusions:** Education (typically completed at young ages) appeared to have a protective effect on lung function later in life, and a small part of this effect was mediated by air pollution.

\* Corresponding author at: Research Department of Epidemiology and Public Health, UCL, London, UK.

E-mail address: [h.pikhart@ucl.ac.uk](mailto:h.pikhart@ucl.ac.uk) (H. Pikhart).

<https://doi.org/10.1016/j.scitotenv.2024.174556>

Received 5 April 2024; Received in revised form 14 June 2024; Accepted 4 July 2024

Available online 5 July 2024

0048-9697/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Studying the associations between outdoor air pollutants and socioeconomic status (SES) is needed as both are important risk factors for numerous health conditions (Glymour et al., 2014; World Health Organization, 2021) including pulmonary outcomes (Orellano et al., 2020; Rocha et al., 2019). Studies in the United States (US) and Europe have reported higher levels of residential exposure to environmental pollutants, particularly PM<sub>2.5</sub> (particulate matter of aerodynamic diameter below 2.5 µm), PM<sub>10</sub> (particulate matter of aerodynamic diameter below 10 µm), and NO<sub>2</sub> (nitrogen dioxide) in people with lower SES (Clark et al., 2014; Hajat et al., 2015; Knobel et al., 2023). However, some European studies found that more deprived groups could be less exposed to air pollutants in urban-rural comparisons, or areas devoted to different economic activities (Fairburn et al., 2019; Padilla et al., 2014; Richardson et al., 2013).

So far, SES's role in the now widely accepted relationship between air pollution and lung function is not fully understood (Hajat et al., 2021). While SES indicators have been proposed to be potential confounders as well as effect modifiers of the association between air pollution and health outcomes (Hajat et al., 2015), it is also possible that air pollution may partly mediate the effects of SES on health (Hajat et al., 2021). Higher educational attainment (as a proxy for SES) may affect the ease (or lack of it) with which individuals choose to live in more (or less) desirable areas (Deluca et al., 2019); these areas might be different in terms of air pollution as well (Wang and Guo, 2023). Furthermore, in as much as educational attainment is related to employment opportunities and income (Glymour et al., 2014), less educated individuals may seek jobs in areas or cities that concentrate polluting industries, thus contributing to higher lifetime air pollution exposures. For example, Padilla et al. (2014) found that the Port of Marseille concentrates poor employees exposed to high levels of NO<sub>2</sub> from steel and petrochemical industries.

SES comprises several dimensions, including income, education, wealth status, and job position; however, education has been among the most widely used in epidemiological research. The vast majority of studies indicate that better-educated individuals fare better concerning health outcomes but also exposure to health risk factors, including air pollution (Cutler and Lleras-Muney, 2010; Donkin, 2014; Hamad et al., 2018). Researchers have investigated the mechanisms underlying the association between education and health, finding evidence for indirect and direct effects. Educational attainment is a powerful predictor of adulthood characteristics, including income and job position, personal control, social support, and health behaviors, which are in turn social determinants of health, indicating an indirect effect on health (Ross and Mirowsky, 2010). Additionally, evidence suggests that individuals with lower SES are more exposed to poorer housing conditions and environmental hazards in their neighborhoods (Evans and Kantrowitz, 2002). Conversely, other researchers argue that education has a more direct impact on health, claiming that it remains a robust predictor of longevity even after controlling for other SES covariates (Baker et al., 2011), and that compulsory schooling laws have shown to be beneficial in reducing mortality, smoking, and obesity (Hamad et al., 2018). Thus, education may influence health both directly and indirectly, serving as a fundamental factor that underlies other SES dimensions.

To our knowledge, the hypothesis that educational attainment drives pulmonary outcomes partly through air pollution exposure (mediator) has not been previously investigated.

Using individual data from a cohort study of Czech older adults from seven small-mid size cities linked with long-term exposures to outdoor air pollutants we aim to investigate (1) the social (educational) gradient of exposure to PM<sub>10</sub> and NO<sub>2</sub> and, (2) the association between exposure to air pollutants and lung functions; (3) the educational gradient in lung function; (4) and the potential mediation effect of air pollution in the association between educational attainment and lung function.

## 2. Methods

### 2.1. Population

We used the baseline survey of the Czech Health, Alcohol, and Psychosocial Factors in Eastern Europe (HAPIEE) cohort study conducted from 2002 to 2005. A population-based urban random sample of older adults was invited to participate in a short physical examination and provide data on health, risk factors, and socioeconomic conditions. Individuals provided their consent to participate. Data were collected by trained personnel. The examination included anthropometric measurements and a spirometry test. A published protocol provides further details (Peasey et al., 2006). A total of 8835 individuals provided demographic and health data, but only 6763 individuals underwent the spirometry test. Of them, 27 had missing values on height and 260 had missing geocodes. Additionally, 119 individuals had missing information on smoking behavior and 47 on education. The analytical sample encompassed 6381 individuals with complete information. Listwise deletion was conducted for missing values as the parameters did not differ from the final sample (Supplement A).

### 2.2. Lung function

Lung function was assessed by trained personnel and standardized protocol using a Micro-Medical Microplus spirometer. The assessment of lung function was conducted in a sitting position on three to eight occasions. The output includes but is not limited to the Forced Vital Capacity (FVC) and the Forced Expiratory Volume in the first second (FEV1). For this study, the maximum value of FEV1 was chosen according to reproducibility and repeatability criteria (Graham et al., 2019). The predicted values of FEV1 were derived using age and height as primary predictors, stratified by sex, using published equations (Kuster et al., 2008). These predicted values were then compared to observed values calculating the ratio of each participant's maximum value of FEV1 to their individual predicted value of FEV1. This ratio was multiplied by 100 for easier interpretation as a percentage difference from the expected FEV1. In consequence, 100 % standardized FEV1 is considered the optimal reference value.

### 2.3. Exposure to air pollution

Air pollution (PM<sub>10</sub> and NO<sub>2</sub>) concentrations between 2005 and 2008 were estimated by the Czech Hydrometeorological Institute (CHMI) on a 1 km × 1 km spatial grid. This estimation employed an empirical inverse distance weighting model, combining several data sources and models: (1) CHMI's Gaussian plume SYMOS'97 atmospheric dispersion model, using the national registry of point and line emission sources (*Information System for Air Quality – ISKO*), traffic-related air pollution data from a 2005 national traffic intensity study, meteorological and digital terrain model data; (2) European Eulerian EMEP (50 km × 50 km) model, to estimate the secondary particle and resuspension concentrations for PM<sub>10</sub>; and (3) altitude data. The models are developed for urban and rural areas separately and merged into spatial maps based on population density grids (Ostatnická, 2008).

The annual means of PM<sub>10</sub> and NO<sub>2</sub> in micrograms per cubic meter (µg/m<sup>3</sup>) were obtained from 2005 to 2008 and matched to the geocoded addresses of the participants. A missing value was assigned if the geographic address could not be retrieved. The 4-year mean concentrations were calculated and used as an approximation to the long-term exposure to PM<sub>10</sub> and NO<sub>2</sub>. Given that interpreting the effect size of small unit changes in PM<sub>10</sub> and NO<sub>2</sub> can be less intuitive, we rescaled both by dividing it by 10. This transformation allows the model coefficients to represent the effect of a 10 µg/m<sup>3</sup> change in PM<sub>10</sub> and NO<sub>2</sub> concentrations, thereby facilitating a clearer understanding of the magnitude of the association of air pollutants and standardized FEV1.

#### 2.4. Socioeconomic status

Participants were asked about their highest level of education achieved. The responses were classified into four categories: (1) primary or incomplete, for those who completed a maximum of 9 years of education; (2) vocational, for those who had four additional years of training in a manual profession; (3) secondary, completed 13 years of education plus a matriculation exam; and (4) tertiary, for individuals with a university degree.

For sensitivity analyses, two additional indicators of SES (deprivation and job as a miner) were used. Three questions were used to assess the level of deprivation: (1) Do you have difficulties in buying food? (2) Do you have difficulties buying clothes? (3) Do you have difficulties paying bills? Answers were formatted on a 4-point Likert scale from 0 (never) to 4 (always). The sum of the three questions ranged from 0 to 12 and were re-categorized as no deprivation (0), low level of deprivation (1–2), and high level of deprivation (3–12). A dummy variable identified whether participants worked as miners (1) or not (0) from free-text occupation descriptions.

#### 2.5. Covariates

Age on the day of the health examination, sex, and smoking behavior were self-reported. Participants were asked if they smoked and had to choose one of the following answers: (1) No, I have never smoked, (2) No, I smoked in the past, but I stopped, (3) Yes, occasionally, less than one cigarette, (4) Yes, regularly, at least one cigarette. These responses were recategorized as follows: (0) never smokers, (1) past and occasional smokers, (2) regular smokers. The body mass index (BMI) was introduced as a continuous variable derived from health examination on weight and height and expressed in  $\text{kg}/\text{m}^2$ .

#### 2.6. Statistical analysis

First, using linear regression, we estimated the association between the level of education and four-year average  $\text{PM}_{10}$  and  $\text{NO}_2$  levels. Multiple linear regressions were used to test the strength of the associations between education and the standardized FEV1 when controlled for covariates in successive models: model 0 was adjusted for age and sex; model 1 additionally adjusted for body mass index (BMI) and smoking status; model 2 was further adjusted for outdoor air pollutants –  $\text{PM}_{10}$  and  $\text{NO}_2$ . The strength of the association between air pollutants and standardized FEV1 was tested using a similar method but model 2 was adjusted for education instead. During the post hoc sensitivity analysis, we included additional indicators of SES (deprivation and job as a miner) to model 2, and to take into account different sources of exposure the results were separated by city category, i.e. presence of mining industry.

Finally, we used structural equation modeling (SEM) builder to perform a mediation analysis. The software output provided the coefficients of the hypothesized paths. We would refer to full mediation if the total direct effect is zero and to partial mediation if it is non-zero. The total direct effect is the coefficient of the path from education to the standardized FEV1. Two indirect effects were calculated: one through  $\text{PM}_{10}$  and one through  $\text{NO}_2$ . To calculate the indirect effect through  $\text{PM}_{10}$ , we multiply the coefficient of education on  $\text{PM}_{10}$  by the coefficient of  $\text{PM}_{10}$  on standardized FEV1. Identical computation was done with the coefficients through  $\text{NO}_2$ . Both indirect effects were summed up to obtain the total indirect effect. To find the percentage mediated we divided the total indirect effect by the sum of total indirect and total direct effects. The model was adjusted for age, sex, smoking, and BMI. The analyses are based on individuals with complete information using Stata v 16.1, at 95 % confidence intervals, and bootstrapping on 5000 resamples for SEM.

### 3. Results

The analytical sample included 6381 individuals, of which 54 % were females (Table 1). The mean age was  $58.24 \pm 7.14$  years. In the period 2005–2008, participants were exposed to a mean of  $33.68 \pm 8.26 \mu\text{g}/\text{m}^3$  of  $\text{PM}_{10}$ , and  $20.57 \pm 9.26 \mu\text{g}/\text{m}^3$  of  $\text{NO}_2$ . Currently, the limit for annual exposure to  $\text{PM}_{10}$  is  $15 \mu\text{g}/\text{m}^3$  and for  $\text{NO}_2$  is  $10 \mu\text{g}/\text{m}^3$  (World Health Organization, 2021). The mean of standardized FEV1 was  $91.91 \pm 17.06$ . Features disaggregated by the level of education showed that tertiary education was more common among males compared to females, higher levels of  $\text{PM}_{10}$  were observed among individuals with lower education levels, and mining cities—Havírov and Karvína—had a higher proportion of individuals with low levels of education.

#### 3.1. Education and air pollution

In age and sex adjusted linear regression analyses,  $\text{PM}_{10}$  exposure was strongly associated with education; specifically, persons with

**Table 1**  
Participants' characteristics by level of educational attainment ( $n = 6381$ ).

	Total	Primary or incomplete	Vocational	Secondary	Tertiary
Age, mean (SD), years	58.24 (7.14)	60.07 (7)	58.15 (7)	57.87 (7)	57.86 (7)
Sex, n (%)					
Male	2917 (46)	157 (6)	1292 (44)	944 (32)	524 (18)
Female	3464 (54)	615 (18)	1083 (31)	1420 (41)	346 (10)
Smoking status, n (%)					
Never	2848 (45)	367 (13)	957 (34)	1073 (38)	451 (16)
Past and occasional	2067 (32)	209 (10)	779 (38)	802 (39)	277 (13)
Regular	1466 (23)	196 (13)	639 (44)	489 (33)	142 (10)
Education					
Primary or incomplete	772 (12)				
Vocational	2375 (37)				
Secondary	2364 (37)				
Tertiary	870 (14)				
BMI, mean (SD), $\text{kg}/\text{m}^2$	28.23 (4.55)	29.60 (5)	28.74 (5)	27.68 (4)	27.13 (4)
Air pollutants, mean (SD), $\mu\text{g}/\text{m}^3$					
$\text{PM}_{10}$ from 2005 to 08	33.68 (8.26)	36.40 (10)	34.54 (9)	32.49 (7)	32.16 (7)
$\text{NO}_2$ from 2005 to 08	20.57 (9.26)	22.15 (9)	20.67 (9)	20.36 (9)	19.42 (10)
<sup>a</sup> Standardized FEV1, mean (SD), % of predicted value	91.91 (17.06)	88.46 (19)	90.87 (17)	92.87 (17)	95.20 (16)
Cities, n (%)					
Havírov	785 (12)	138 (18)	357 (45)	214 (27)	76 (10)
Hradec	732 (11)	50 (7)	235 (32)	315 (43)	132 (18)
Jihlava	657 (10)	78 (12)	264 (40)	244 (37)	71 (11)
Karvína	478 (7)	116 (24)	230 (48)	107 (23)	25 (5)
Kromeriz	1271 (20)	118 (9)	434 (34)	492 (49)	227 (18)
Liberec	876 (14)	70 (8)	287 (33)	377 (43)	142 (16)
Usti nad Labem	1582 (25)	202 (13)	568 (36)	615 (39)	197 (12)

<sup>a</sup> Optimal value of standardized FEV1 = 100 %.

tertiary education had significantly lower exposures than those with primary or incomplete education, although the difference was small in absolute terms, about 4  $\mu\text{g}/\text{m}^3$  (Fig. 1, left panel). A similar relationship was observed between  $\text{NO}_2$  and education: people at the highest level of education had lower exposure to  $\text{NO}_2$  (Fig. 1, right panel).

### 3.2. Determinants of lung function

The simultaneous associations of  $\text{PM}_{10}$  and  $\text{NO}_2$  with the standardized FEV1 adjusted for multiple risk factors in ordered steps are presented in Table 2. When adjusted for age and sex (model 0), a 10  $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{10}$  was associated with 1.78 % (95 % CI: -2.33 to -1.23) lower FEV1, while a 10  $\mu\text{g}/\text{m}^3$  increase in  $\text{NO}_2$  was not significantly related to FEV1 (0.29, 95 % CI: -0.20 to 0.78). The introduction of BMI and smoking (model 1) and then education (model 2) explained part of the association between  $\text{PM}_{10}$  and FEV1. No changes in the association between  $\text{NO}_2$  and standardized FEV1 were observed with the addition of covariates.

After adjusting for multiple factors in successive linear regressions, a strong educational gradient was observed as individuals with higher levels of education had better lung function than those in an immediately inferior category (Table 3). In model 0, the regression coefficients showed that people with tertiary education had seven percentage points higher standardized FEV1 than people with primary or incomplete education. When BMI and smoking were included (model 1) in the regression analysis, the differences between the effect of the highest and the lowest education categories on FEV1 became smaller. However, there remained almost a five-point gap in standardized FEV1 between the lowest and highest education groups. Model 2, with air pollutants added, reduced the educational gradient by only an extra 0.51 points in standardized FEV1 compared to model 1.

In sensitivity analyses, first, we tested for interactions between sex and level of education and found non-significant differences (Supplement B). Then, given the differences in educational attainment between Havirov and Karvina, cities historically devoted to the mining industry, compared to other cities (Supplement C), a separate analysis by city category of model 2 was considered (Supplement D). Finally, we tested the strength of the associations when other SES indicators were included. Table 4 presents the regression coefficients for the association of educational attainment, deprivation, job as a miner, and air pollutants with standardized FEV1. Education had a strong positive association with standardized FEV1 in both groups of cities. In Havirov and Karvina, people with tertiary education had almost six percentage points higher standardized FEV1 than people with primary or incomplete education.

**Table 2**

Linear regression coefficients for the association of  $\text{PM}_{10}$  and  $\text{NO}_2$  with standardized FEV1(%) among Czech adults participating in the HAPPIE study.

	Model 0	Model 1	Model 2
	b <sup>a</sup> (95 % CI)	b <sup>a</sup> (95 % CI)	b <sup>a</sup> (95 % CI)
$\text{PM}_{10}$ per 10 $\mu\text{g}/\text{m}^3$	-1.78 (-2.33 to -1.23)	-1.53 (-2.07 to -0.99)	-1.33 (-1.87 to -0.78)
$\text{NO}_2$ per 10 $\mu\text{g}/\text{m}^3$	0.29 (-0.20 to 0.78)	0.35 (-0.12 to 0.83)	0.36 (-0.12 to 0.84)
AIC	54,227.07	53,909.11	53,884.48
BIC	54,260.88	53,963.21	53,958.85
LR test		$\chi^2$ (3) =323.95, p < 0.001	$\chi^2$ (3) =30.64, p < 0.001

Model 0: adjusted for age and sex.

Model 1: as model 0 additionally adjusted for BMI and smoking status.

Model 2: as model 1 additionally adjusted for education.

<sup>a</sup> b coefficient of -1 means that standardized FEV1 reduces by one percentage point per 10  $\mu\text{g}/\text{m}^3$  increase in air pollutant concentration.

**Table 3**

Linear association of educational attainment with standardized FEV1(%) among Czech adults participating in the HAPPIE study.


	Model 0	Model 1	Model 2
	b <sup>a</sup> (95 % CI)	b <sup>a</sup> (95 % CI)	b <sup>a</sup> (95 % CI)
Primary or incomplete	0	0	0
Vocational	3.00 (1.59 to 4.41)	2.41 (1.03 to 3.79)	2.17 (0.78 to 3.55)
Secondary	4.61 (3.22 to 6.00)	3.19 (1.82 to 4.57)	2.71 (1.33 to 4.09)
Tertiary	7.43 (5.76 to 9.11)	5.16 (3.49 to 6.82)	4.65 (2.97 to 6.32)
AIC	54,184.68	53,903.66	53,884.47
BIC	54,225.24	53,964.51	53,958.84
LR test		$\chi^2$ (3) =287.01, p < 0.001	$\chi^2$ (2) =23.19, p < 0.001

Model 0: adjusted for age and sex.

Model 1: as model 0 additionally adjusted for BMI and smoking status.

Model 2: as model 1 additionally adjusted for  $\text{PM}_{10}$  and  $\text{NO}_2$ .

<sup>a</sup> b coefficient of -1 means that standardized FEV1 reduces by one percentage point per category of education compared to individuals with primary or incomplete education.



**Fig. 1.** Linear regression coefficients of categories of attained education for exposure to  $\text{PM}_{10}$  and  $\text{NO}_2$ . Coefficients and confidence intervals are adjusted for age and sex.



**Table 4**

Multivariable linear association coefficients of variables included in model 2 plus other SES indicators (deprivation and job as miner) on Standardized FEV1.  $n = 6187$ .

	Mining cities (Havirov and Karvina)		Non-mining cities (Hradec, Jihlava, Kromeriz, Liberec, Usti nad Labem)	
	b	95 % CI	b	95 % CI
<b>Education</b>				
Primary or incomplete	Ref.		Ref.	
Vocational	2.88	0.23 to 5.53	1.40	-0.30 to 3.11
Secondary	4.48	1.60 to 7.35	1.81	0.13 to 3.48
Tertiary	5.65	1.54 to 9.76	3.69	1.72 to 5.65
<b>Air pollutants</b>				
<sup>a</sup> PM <sub>10</sub> per 10 µg/m <sup>3</sup>	-4.98	-8.83 to -1.14	-1.16	-2.60 to 0.28
<sup>a</sup> NO <sub>2</sub> per 10 µg/m <sup>3</sup>	1.08	-0.70 to 2.85	-0.28	-0.23 to 0.79
<b>Deprivation</b>				
No deprivation	Ref.		Ref.	
Low deprivation	-1.84	-4.17 to 0.49	-0.80	-1.93 to 0.33
High deprivation	-2.49	-4.73 to -0.24	-1.11	-2.27 to 0.06
Miner, yes	-2.99	-5.90 to -0.08		

<sup>a</sup> Average concentration of air pollutants from 2005 to 2008 expressed in µg/m<sup>3</sup>. b coefficient of -1 means that standardized FEV1 reduces by one percentage point per 10 µg/m<sup>3</sup> increase in air pollutant concentration.

PM<sub>10</sub> had a stronger negative effect in the historically mining cities of Havirov and Karvina: a 10 µg/m<sup>3</sup> increase in PM<sub>10</sub> reduced the standardized FEV1 by five points, while in the non-mining cities, the reduction was not statistically significant. NO<sub>2</sub> showed no significant associations with standardized FEV1. Additionally, in Havirov and Karvina people with the highest level of deprivation showed a 2.49 % reduction in standardized FEV1 and miners lost 2.99 % in standardized FEV1 (Table 4). Deprivation was not associated with lower lung function in non-mining cities.

3.3. Mediation analysis

Table 5 and Fig. 2 display the effects of the proposed paths. All paths show statistically significant associations, except for the path from NO<sub>2</sub> to standardized FEV1. Higher educational attainment was associated with less exposure to PM<sub>10</sub> and less exposure to NO<sub>2</sub>. PM<sub>10</sub> was negatively associated with the standardized FEV1, which means that higher levels of air pollution led to poorer lung function. The increase in 10 µg/m<sup>3</sup> of PM<sub>10</sub> was associated with reduced standardized FEV1 by 1.32 %.


The coefficients from education were multiplied by the coefficients toward standardized FEV1 to obtain the indirect effects specific to each mediator (Table 6). We found significant indirect effects through PM<sub>10</sub> (b = 0.20, CI 95 %: 0.11 to 0.29) but not for NO<sub>2</sub> (b = -0.03, -0.06 to 0.01). The total indirect effect through PM<sub>10</sub> and NO<sub>2</sub> accounts for 12 %

**Table 5**

Mediation effects of PM<sub>10</sub> and NO<sub>2</sub> in the relationship between educational attainment and standardized FEV1(%) among Czech adults participating in the HAPPIE study.

Exposure	Outcome	b	Standard error	95 % CI
	Standardized FEV1 (%)	1.34	0.250	0.85 to 1.83
Education	PM <sub>10</sub> , µg/m <sup>3</sup>	-0.15	0.011	-0.18 to -0.13
	NO <sub>2</sub> , µg/m <sup>3</sup>	-0.07	0.013	-0.10 to -0.05
PM <sub>10</sub> per 10 µg/m <sup>3</sup>	Standardized FEV1 (%)	-1.32	0.283	-1.87 to -0.76
NO <sub>2</sub> per 10 µg/m <sup>3</sup>		0.34	0.246	-0.14 to 0.83

Model adjusted for age, sex, BMI, and smoking behavior.



**Fig. 2.** Mediation effects of PM<sub>10</sub> and NO<sub>2</sub> in relationship between education and standardized FEV1(%). \*Statistically significant coefficients. Model adjusted for age, sex, BMI, and smoking behavior. Model fit: X<sup>2</sup>(9) = 177.337,  $p < 0.001$ , CFI = 0.775, RMSEA = 0.054, 90 % CI RMSEA [0.047, 0.061].

of the total effect.

4. Discussion

In this Czech population-based urban study, most of our findings were consistent with the existing literature. First, exposure to higher levels of PM<sub>10</sub> was associated with worse lung function (Doiron et al., 2019), and we confirmed educational gradients in exposure to air pollutants and lung function. Given these observations, we explored the potential mediation effect of air pollution between educational attainment and lung function. We found a partial mediation: around 12 % of the total effect of educational attainment on lung function can be attributed to exposure to air pollution. This finding supports our hypothesis that the level of education could influence the degree of air pollution people are exposed to. Nevertheless, 88 % of the effect of educational attainment on lung function remained unexplained by air pollutants.

The educational gradient observed for air pollutants, for FEV1, and the partial mediation detected suggest that the level of attained education may at least partly determine exposure to outdoor air pollution later in life. The highest level of education is generally completed by early adulthood, thus other characteristics of SES could better explain the association with air pollution exposure and lung function in late adulthood. However, in this sample of Czech individuals, education was a stronger socioeconomic predictor of long-term health outcomes. Furthermore, we have already described the association between social mobility and lung function elsewhere (Quispe-Haro et al., 2022). Given the cross-sectional design and lack of additional information, we were neither able to quantify the residential relocations during the life course of HAPPIE study participants, nor to test whether being born in a polluted or deprived area could lead to lower levels of attained education in our sample. Others have explored the associations between the migration of talented individuals, exposure to air pollutants, and socioeconomic conditions (Germani et al., 2021; Wang and Guo, 2023). The reasons for voluntary or involuntary migration are complex and diverse. Individuals with any level of education migrate and consider not only housing quality and job opportunities but also family ties, costs, and familiarity (Deluca et al., 2019); hence longitudinal data is needed to analyze to what extent people consider the level of air pollution in the residential area and the posterior health consequences when relocating.

Epidemiologists have focused on the effect modification by SES in the relationship between air pollution and pulmonary outcomes. A systematic review found weak evidence of effect modification of SES when evaluating the association between air pollution and asthma

**Table 6**

Bootstrapped indirect effects coefficients in the mediation analysis between education and standardized FEV1(%).

	b	Standard error	95 % Confidence interval
Total indirect effect	0.18	0.043	0.09 to 0.26
PM <sub>10</sub> , µg/m <sup>3</sup>	0.20	0.046	0.11 to 0.29
NO <sub>2</sub> , µg/m <sup>3</sup>	-0.03	0.019	-0.06 to 0.01

exacerbation in children (Rodriguez-Villamizar et al., 2016). For low-income adults, stronger harmful effects of air pollution on FEV1 and FVC were reported, when controlled by the level of education (Doiron et al., 2019). Baker et al. (2011) argue that the causal role of education on health has been questioned in part due to the lack of sophisticated statistical approaches to “control for wealth and other dimensions of socioeconomic status”. In our sample, the educational gradient remained strong after stratifying the analysis between mining and non-mining cities and adding other SES indicators alongside education, such as working as a miner and level of economic deprivation. In additional analyses, we rejected other probable mechanisms involved, including the interaction between education and sex explored in other studies (Forastiere et al., 2007; Hajat et al., 2021).

As proposed by Hajat et al. (2021), economic constraints could force people of lower SES to live near roads or industrial areas where housing prices are affordable. However, in Europe geographic heterogeneity of social conditions and air pollution have been described (Padilla et al., 2014; Stroh et al., 2005), suggesting that the environmental inequalities found might be specific for the studied area. In our data, low-educated individuals were exposed to higher levels of both PM<sub>10</sub> and NO<sub>2</sub>. However, these results are inconsistent with previous ecological observations of the distribution of air pollutants in the Czech Republic (Branis and Linhartova, 2012), where higher levels of NO<sub>2</sub> were associated with a higher proportion of citizens with university education. This may be explained, firstly, by the sample consisting solely of individuals living in urban areas and the vicinity of major road networks, generally predicting transport accessibility and mobility, that can be associated with higher SES scores, especially when compared to peripheral areas. Additionally, the discrepancy can be attributed to the spatial resolution used, which was 1 × 1 km, potentially leading to less precise estimates. Compared to PM<sub>10</sub>, NO<sub>2</sub> has different distance-decay gradients, a predominant association with line sources (traffic-related air pollution), and thus higher variance within the grid cells associated more with area/volume sources with less spatial variability (McAdam et al., 2011). Thus, unlike PM<sub>10</sub>, the NO<sub>2</sub> levels may instead indicate the urban economic centrality. Inaccuracies in the computation of air pollution exposure can lead to exposure misclassification (Lu, 2023). Consequently, if more precise estimates of air pollution were used, the proportion mediated by air pollutants (12 %), though significant, might be slightly different. This is primarily because the effect size was driven by PM<sub>10</sub> rather than NO<sub>2</sub>. We speculate that a more spatially precise measurement of NO<sub>2</sub> could help improve our models.

Most evidence on the association of SES, air pollution exposure, and health outcomes comes from Western countries where income or job position could be more appropriate to describe their social standing. Western countries with high educational attainment have a limited variation in schooling which could explain the limited capacity of education to capture the health effects (Baker et al., 2011). On the contrary, the educational attainment in the Czech Republic might be a better indicator of the social hierarchy. Previous analyses of the HAPIEE study have demonstrated that education is the strongest socioeconomic predictor of health outcomes (Vandenheede et al., 2014),

This study had several limitations that must be acknowledged. Firstly, a single indicator of SES was used in our mediation analysis, but other aspects of SES could independently predict health outcomes, such as control over one's own life and job, income, and the self-perception of position in the social hierarchy (Marmot, 2004; Theodossiou and Zangelidis, 2009). Moreover, the study did not account for the influence of time spent outdoors and commuting (Makri and Stilianakis, 2008), or the contribution of indoor air pollution on total exposure to air pollutants, which could have potential implications for the results. The cross-sectional nature of the data does not allow us to assess temporality, although – at least in relation to education – reverse causality is unlikely. Nevertheless, a long-term follow-up study would provide more robust evidence.

Air pollution levels have improved in Western Europe in general as

well as in the Czech Republic since the early 2000 (Shen et al., 2022), hence it is probable that participants' lifetime exposure to outdoor pollution was substantially higher, therefore we could underestimate the contribution of air pollutants. The lack of consideration for residential mobility further adds to the study's limitations. We assumed that the level of education would affect adults in terms of job opportunities and mobility, however, these speculations should be supported by evidence that will consider the conscious or unconscious decisions of people in terms of career, income, family, housing and environmental expectations, and opportunities.

To address these issues and enhance future research, incorporating critical concepts such as the social exposome perspective, life course analysis, and differential vulnerability would be valuable (Gudi-Mindermann et al., 2023). Additionally, considering time spent in various activities that influence exposure outside residence can provide more comprehensive insights into the intricate interactions between air pollutants, health effects, and socioeconomic factors. By acknowledging and rectifying these limitations, future studies can contribute significantly to the understanding and management of this crucial public health issue.

## 5. Conclusion

Individuals with lower educational attainment (representing lower SES) are likely to be exposed to worse air quality. We found evidence of a partial mediation effect of air pollution in the association of education and lung function. Higher levels of education showed both a protective direct effect on lung function and a protective effect on exposure to air pollutants. Higher spatial resolution of NO<sub>2</sub> concentration data could help to capture the effects on lung function. Insights from other fields should be incorporated to understand better the relationship between an individual's socioeconomic circumstances, mobility patterns, environmental exposures, and health.

## CRedit authorship contribution statement

**Consuelo Quispe-Haro:** Writing – original draft, Visualization, Methodology, Formal analysis, Conceptualization. **Daniel Szabó:** Data curation. **Katarzyna Kordas:** Writing – review & editing, Methodology. **Nadezda Capkova:** Writing – review & editing. **Hynek Pikhart:** Writing – review & editing, Supervision, Methodology. **Martin Bobak:** Writing – review & editing, Supervision, Methodology.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Data availability

Data will be made available on request.

## Acknowledgment

Authors thank the Research Infrastructure RECETOX (number LM2023069) and project CETOCOEN EXCELLENCE (number CZ.02.1.01/0.0/0.0/17\_043/0009632) financed by the Czech Ministry of Education, Youth and Sports for supportive background. This work was supported by the European Union's Horizon 2020 Research and Innovation Programme under grant agreement number 857560 (CETOCOEN Excellence) and 857487 (R-Exposome Chair). This publication reflects only the author's view, and the European Commission is not responsible for any use that may be made of the information it contains. KK was supported by the 2023-24 Czech Fullbright Commission Distinguished Scholar Award.

## Funding

The authors have disclosed the receipt of the following financial support for the research, authorship and/or publication of this article: the HAPIEE study was funded by the Wellcome Trust (grant WT064947 and WT081081), the US National Institute of Aging (grant R01 AG23522) and the MacArthur Foundation. This work was financially supported by the project Systemic Risk Institute (LX22NPO5101, NPO - EXCELES) funded by the European Union - Next Generation EU.

## Ethics approval

This study involves human participants and was approved by the University College London, UK and by the local ethics committee in every participating centre. Participants gave informed consent to participate in the study before taking part. The study protocols were approved by ethical committees at University College London, UK, and at each participating centre.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2024.174556>.

## References

- Baker, D.P., Leon, J., Smith Greenaway, E.G., Collins, J., Movit, M., 2011. The education effect on population health: a reassessment. *Popul. Dev. Rev.* 37, 307–332. <https://doi.org/10.1111/j.1728-4457.2011.00412.x>.
- Branis, M., Linhartova, M., 2012. Association between unemployment, income, education level, population size and air pollution in Czech cities: evidence for environmental inequality? A pilot national scale analysis. *Health Place* 18, 1110–1114. <https://doi.org/10.1016/j.healthplace.2012.04.011>.
- Clark, L.P., Millet, D.B., Marshall, J.D., 2014. National patterns in environmental injustice and inequality: outdoor NO<sub>2</sub> air pollution in the United States. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0094431>.
- Cutler, D.M., Lleras-Muney, A., 2010. Understanding differences in health behaviors by education. *J. Health Econ.* 29, 1–28. <https://doi.org/10.1016/j.jhealeco.2009.10.003>.
- Deluca, S., Wood, H., Rosenblatt, P., 2019. Why poor people move (and where they go): residential mobility, selection and stratification. *City Community* 18, 556–593. <https://doi.org/10.1111/cico.12386>.
- Doiron, D., de Hoogh, K., Probst-Hensch, N., Fortier, I., Cai, Y., de Matteis, S., Hansell, A. L., 2019. Air pollution, lung function and COPD: results from the population-based UK Biobank study. *Eur. Respir. J.* 54 <https://doi.org/10.1183/13993003.02140-2018>.
- Donkin, A.J.M., 2014. Social Gradient. In: Cockerham, W.C., Dingwall, R., Quah, S. (Eds.), *The Wiley Blackwell Encyclopedia of Health, Illness, Behavior, and Society*. John Wiley & Sons, Chichester, pp. 2172–2178. <https://doi.org/10.1002/9781118410868.wbehubs530>.
- Evans, G.W., Kantrowitz, E., 2002. Socioeconomic status and health: the potential role of environmental risk exposure. *Annu. Rev. Public Health.* <https://doi.org/10.1146/annurev.publhealth.23.112001.112349>.
- Fairburn, J., Schüle, S.A., Dreger, S., Hilt, L.K., Bolte, G., 2019. Social inequalities in exposure to ambient air pollution: a systematic review in the WHO European region. *Int. J. Environ. Res. Public Health* 16. <https://doi.org/10.3390/ijerph16173127>.
- Forastiere, F., Stafoggia, M., Tasco, C., Picciotto, S., Agabiti, N., Cesaroni, G., Perucci, C. A., 2007. Socioeconomic status, particulate air pollution, and daily mortality: differential exposure or differential susceptibility. *Am. J. Ind. Med.* 50, 208–216. <https://doi.org/10.1002/ajim.20368>.
- Germani, A.R., Scaramozzino, P., Castaldo, A., Talamo, G., 2021. Does air pollution influence internal migration? An empirical investigation on Italian provinces. *Environ. Sci. Policy* 120, 11–20. <https://doi.org/10.1016/j.envsci.2021.02.005>.
- Glymour, M., Avendano, M., Kawachi, I., 2014. Socioeconomic status and health. In: Brakman, L., Kawachi, I., Glymour, M. (Eds.), *Social Epidemiology*. Oxford University Press, New York, pp. 17–62.
- Graham, B.L., Steenbruggen, I., Barjaktarevic, I.Z., Cooper, B.G., Hall, G.L., Hallstrand, T. S., Kaminsky, D.A., McCarthy, K., McCormack, M.C., Miller, M.R., Oropez, C.E., Rosenfeld, M., Stanojevic, S., Swanney, M.P., Thompson, B.R., 2019. Standardization of spirometry 2019 update. An Official American Thoracic Society and European Respiratory Society Technical Statement. <https://doi.org/10.1164/rccm.201908-1590ST>, 200, E70–E88.
- Gudi-Mindermann, H., White, M., Roczen, J., Riedel, N., Dreger, S., Bolte, G., 2023. Integrating the social environment with an equity perspective into the exposome paradigm: a new conceptual framework of the social Exposome. *Environ. Res.* 233, 116485 <https://doi.org/10.1016/j.envres.2023.116485>.
- Hajat, A., Hsia, C., O'Neill, M.S., 2015. Socioeconomic disparities and air pollution exposure: a global review. *Curr. Environ. Health Rep.* <https://doi.org/10.1007/s40572-015-0069-5>.
- Hajat, A., Maclehorse, R.F., Rosofsky, A., Walker, K.D., Clougherty, J.E., 2021. Confounding by socioeconomic status in epidemiological studies of air pollution and health: challenges and opportunities. *Environ. Health Perspect.* 129 <https://doi.org/10.1289/EHP7980>.
- Hamad, R., Elser, H., Tran, D.C., Rehkopf, D.H., Goodman, S.N., 2018. How and why studies disagree about the effects of education on health: a systematic review and meta-analysis of studies of compulsory schooling laws. *Soc. Sci. Med.* <https://doi.org/10.1016/j.socscimed.2018.07.016>.
- Knobel, P., Hwang, I., Castro, E., Sheffield, P., Holaday, L., Shi, L., Amini, H., Schwartz, J., Yitshak Sade, M., 2023. Socioeconomic and racial disparities in source-apportioned PM<sub>2.5</sub> levels across urban areas in the contiguous US, 2010. *Atmos. Environ.* 303. <https://doi.org/10.1016/j.atmosenv.2023.119753>.
- Kuster, S.P., Kuster, D., Schindler, C., Rochat, M.K., Braun, J., Held, L., Brändli, O., 2008. Reference equations for lung function screening of healthy never-smoking adults aged 18–80 years. *Eur. Respir. J.* 31, 860–868. <https://doi.org/10.1183/09031936.00091407>.
- Lu, Y., 2023. Assessing air pollution exposure misclassification using high-resolution PM<sub>2.5</sub> concentration model and human mobility data. *Air Qual. Atmos. Health* 16, 2225–2238. <https://doi.org/10.1007/s11869-023-01404-2>.
- Makri, A., Stilianakis, N.I., 2008. Vulnerability to air pollution health effects. *Int. J. Hyg. Environ. Health* 211, 326–336. <https://doi.org/10.1016/j.ijheh.2007.06.005>.
- Marmot, M., 2004. Status syndrome. *Significance* 1, 150–154. <https://doi.org/10.1111/j.1740-9713.2004.00058.x>.
- McAdam, K., Steer, P., Perrotta, K., 2011. Using continuous sampling to examine the distribution of traffic related air pollution in proximity to a major road. *Atmos. Environ.* 45, 2080–2086. <https://doi.org/10.1016/j.atmosenv.2011.01.050>.
- Orellano, P., Reynoso, J., Quaranta, N., Bardach, A., Ciapponi, A., 2020. Short-term exposure to particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>), and ozone (O<sub>3</sub>) and all-cause and cause-specific mortality: systematic review and meta-analysis. *Environ. Int.* 142 <https://doi.org/10.1016/j.envint.2020.105876>.
- Ostatnická, J., 2008. Znečištění ovzduší na území České republiky v roce 2008 [WWW Document]. <https://www.chmi.cz/files/portal/docs/uoco/isko/grafroc/groc/gr08cz/kap22.html> (accessed 3.5.24).
- Padilla, C.M., Kihal-Talantikite, W., Vieira, V.M., Rossello, P., Nir, Le, G., Zmirou-Navier, D., Deguen, S., 2014. Air quality and social deprivation in four French metropolitan areas—a localized spatio-temporal environmental inequality analysis. *Environ. Res.* 134, 315–324. <https://doi.org/10.1016/j.envres.2014.07.017>.
- Peasey, A., Bobak, M., Kubinova, R., Malyutina, S., Pajak, A., Tamosiunas, A., Pikhart, H., Nicholson, A., Marmot, M., 2006. Determinants of cardiovascular disease and other non-communicable diseases in central and Eastern Europe: rationale and design of the HAPIEE study. *BMC Public Health* 6, 1–10. <https://doi.org/10.1186/1471-2458-6-255/TABLES/2>.
- Quispe-Haro, C., Pajak, A., Tamosiunas, A., Capkova, N., Bobak, M., Pikhart, H., 2022. Socioeconomic position over the life course and impaired lung function of older adults in central and Eastern Europe: the HAPIEE study. *J. Epidemiol. Community Health* 1978 (77), 49–55. <https://doi.org/10.1136/jech-2022-219348>.
- Richardson, E.A., Pearce, J., Tunstall, H., Mitchell, R., Shortt, N.K., 2013. Particulate air pollution and health inequalities: a Europe-wide ecological analysis. *Int. J. Health Geogr.* 12 <https://doi.org/10.1186/1476-072X-12-34>.
- Rocha, V., Soares, S., Stringhini, S., Fraga, S., 2019. Socioeconomic circumstances and respiratory function from childhood to early adulthood: a systematic review and meta-analysis. *BMJ Open* 9. <https://doi.org/10.1136/bmjopen-2018-027528>.
- Rodriguez-Villamizar, L.A., Berney, C., Villa-Roel, C., Ospina, M.B., Osornio-Vargas, Alvaro, Rowe, B.H., 2016. The role of socioeconomic position as an effect-modifier of the association between outdoor air pollution and children's asthma exacerbations: an equity-focused systematic review. *Rev. Environ. Health* 31, 297–309.
- Ross, C., Mirowsky, J., 2010. Why education is the key to socioeconomic differentials in health. In: Bird, C., Conrad, P., Fremont, A., Timmermans, S. (Eds.), *Handbook of Medical Sociology*. Vanderbilt University Press, Nashville, pp. 33–51.
- Shen, Y., de Hoogh, K., Schmitz, O., Clinton, N., Tuxen-Bettman, K., Brandt, J., Christensen, J.H., Frohn, L.M., Geels, C., Karssenberg, D., Vermeulen, R., Hoek, G., 2022. Europe-wide air pollution modeling from 2000 to 2019 using geographically weighted regression. *Environ. Int.* 168, 107485 <https://doi.org/10.1016/j.envint.2022.107485>.
- Stroh, E., Oudin, A., Gustafsson, S., Pilesjö, P., Harrie, L., Strömberg, U., Jakobsson, K., 2005. Are associations between socio-economic characteristics and exposure to air pollution a question of study area size? An example from Scania, Sweden. *Int. J. Health Geogr.* 4 <https://doi.org/10.1186/1476-072X-4-30>.
- Theodossiou, I., Zangelidis, A., 2009. The social gradient in health: the effect of absolute income and subjective social status assessment on the individual's health in Europe. *Econ. Hum. Biol.* 7, 229–237. <https://doi.org/10.1016/j.ehb.2009.05.001>.

- Vandenheede, H., Vikhireva, O., Pikhart, H., Kubinova, R., Malyutina, S., Pajak, A., Tamosiunas, A., Peasey, A., Simonova, G., Topor-Madry, R., Marmot, M., Bobak, M., 2014. Socioeconomic inequalities in all-cause mortality in the Czech Republic, Russia, Poland and Lithuania in the 2000s: findings from the HAPIEE study. *J. Epidemiol. Community Health* 1978 (68), 297–303. <https://doi.org/10.1136/jech-2013-203057>.
- Wang, H., Guo, F., 2023. City-level socioeconomic divergence, air pollution differentials and internal migration in China: migrants vs talent migrants. *Cities* 133. <https://doi.org/10.1016/j.cities.2022.104116>.
- World Health Organization, 2021. WHO Global Air Quality Guidelines. Particulate Matter (PM2.5 and PM10), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide. Geneva.